arXiv:1602.03375v2 [quant-ph] 25 Feb 2016

Two new integrable cases of two-dimensional quantum mechanics with a magnetic field

V.G. Marikhin *
L.D. Landau Institute for Theoretical Physics, RAS
Chernogolovka, Moscow region, Russia

February 26, 2016

Abstract

Two integrable cases of two-dimensional Schrödinger equation with a magnetic field are proposed. Using the polar coordinates and the symmetrical gauge, we will obtain solutions of these equation through Biconfluent and Confluent Heun functions. The quantization rules will be derived for both systems under consideration.

Key words: quantum mechanics, magnetic field, special functions
PACS numbers: 03.65.-w, 02.30.Gp

The problem of obtaining integrable cases of the Schrödinger equation has a long history. The number of such cases is not very large. The most famous examples can be found in [1], see also the celebrated work by Infeld and Hull [2] and the classic papers by Schrödinger himself [3]–[5]. Among the recent studies we can highlight [6]–[9]. Most of these works deal with one-dimensional problems.

The problem of integrable cases of the two-dimensional Schrodinger equation with an electromagnetic field has long been studied as well, including the celebrated "Landau levels" [1], (see e.g. [10] and the references therein), but it is not completely solved yet. After the appearance of modern methods for analyzing integrable systems, various aspects of this problem were investigated. For instance, the class of (finite-gap) solutions of the Schrödinger equation with a magnetic field was studied in [11] (see also [12]). The factorization method was used in [10] to solve this problem (see also the references therein). The Schrödinger equation in a magnetic field with additional linear and quadratic integrals of motion were considered, for instance, in [13] (also see the references therein), where several interesting examples of such equations in different coordinate systems were obtained.

The goal of this article is to introduce two new integrable examples of the two-dimensional Schrödinger equation with an electromagnetic field. We integrate these cases and describe the structure of the quantum states in terms of the Biconfluent and Confluent Heun functions. We consider only the discrete spectrum and the wave functions of the corresponding Schrödinger operator.

*e-mail: mvg@itp.ac.ru

We need some properties of the above mentioned Biconfluent and Confluent Heun functions, especially, the conditions when these functions reduce to polynomials. One can find the necessary information in many works [14]–[19], but for the reader's convenience and for further applications, we re-obtain some of these properties in Appendix.

The author's work [20] contains a classification of all two-dimensional Schrödinger operators with an additional integral of motion quadratic in the momenta. Shortly speaking, one can derive new examples by starting from some solution of the quasi-Stäckel system (in Stäckel coordinates) and transforming this solution into the usual Cartesian coordinates. However, it is a non-trivial problem to distinguish the examples which are interesting from the physical standpoint, because of the large number of parameters contained in the classification results. We postpone the comprehensive analysis of this problem for the future work. At the moment, we present just two particular examples with a reduced set of parameters. Namely, we consider the Schrödinger operator in the usual form

$$\hat{H} = \frac{1}{2m}\left(-i\hbar\vec{\nabla} - \frac{e}{c}\vec{A}\right)^2 + u(r,\phi), \quad \hat{H}\psi = E\psi, \tag{1}$$

where the vector potential A and the potential u presented below contain three independent parameters: the length a, a dimensionless parameter ϵ and parameter k which is integer in the case of the discrete spectrum.

Example 1. Repulsive potential

Let us consider the example defined by the vector potential and potential

$$A_\phi = \frac{c\hbar}{2e\,a^2}\, r\left(\epsilon + 3\frac{r^2}{a^2}\right), \quad A_r = 0, \quad u = -\frac{\hbar^2}{2ma^2}\left(2\frac{r^6}{a^6} + \epsilon\frac{r^4}{a^4} + 2k\frac{r^2}{a^2}\right), \tag{2}$$

and the energy

$$E = \frac{\hbar^2}{2ma^2}\lambda. \tag{3}$$

We calculate a magnetic field

$$B = \frac{c\hbar}{ea^2}(\epsilon + 6\frac{r^2}{a^2}).$$

At a first glance, the situation looks dangerous, because the potential in (2) is strongly repulsive. However, it turns out that the contribution of the magnetic field prevails the contribution of the potential. A method to obtain an appropriate wave function is the following.

1. We substitute into (1) the wave function of the form

$$\psi = e^{-\frac{r^4}{8a^4}} e^{-\epsilon\frac{r^2}{4a^2}} r^l e^{il\phi} w(r), \tag{4}$$

where $l \geq 0$. The equation for w is solved in terms the Biconfluent Heun function

$$w(r) = \mathrm{HeunB}(l, \epsilon, 3l + 2k, -\epsilon l - \lambda, \frac{r^2}{2a^2}) \tag{5}$$

with the parameters $\alpha = l,\ \beta = \epsilon,\ \gamma = 3l + 2k, \delta = -l\epsilon - \lambda$, while its second linearly independent solution is irregular in the point $r = 0$. The requirement of the regularity of the

HeunB function (5) in the whole range of definition implies that it must be a polynomial (see [18] and citations therein for more details). The polynomiality conditions are given in Appendix.

The condition (23) in this case becomes $l+k=n+1$. Thus, $k \leq n+1$. We note that in this example $\det(A_{n+1})$ is a polynomial in λ of degree $n+1$ and the condition $\det(A_{n+1})=0$ defines $n+1$ values λ^i $(i=1,2,...,n+1)$ and therefore $n+1$ energy levels E^i by formula (3).

To obtain the corresponding wave function, we substitute $\lambda=\lambda_i$ into the matrix A_{n+1}, compute $n+1$ eigenvectors $\vec{p^i}=(p_0^i,p_1^i,..p_n^i)$ of this matrix and obtain the polynomials $w^i(r)=P_n^i(\frac{r^2}{2a^2})$, where $P_n^i=\sum\limits_{j=0}^{n} p_j^i z^j$. Finally, we substitute $w^i(r)$ into formula (4) in order to obtain $n+1$ wave functions ψ^i.

At the moment, we have described the quantum states at fixed n (or fixed $l=n+1-k$). In order to obtain the full picture, we have to fix the Hamiltonian parameter k, to consider all possible non-negative integers l or $n=k-1,k,k+1,k+2,...$ and to obtain an infinite set of "blocks" which contain $n+1$ energy levels and wave functions for each permissible $n \geq k-1$. Notice, that i is an "internal" index for each n.

It is very important that only real roots λ of equation (24) in this example are physical and belong to the discrete spectrum as well as corresponding energies E and the wave functions ψ. Therefore we must choose only these roots.

Thus, we obtain a rather complicated structure of the quantum (discrete) states, which is non-typical for the quantum mechanics. Such structure is similar for all cases under consideration.

All wave functions (4) under consideration are strongly localized in the length scale of a by the quadratic on r magnetic field in this example.

We note that the diagonals of the matrix A_{n+1} (22) in this example are defined by the following sequences:

$$a_j=\lambda-\epsilon(2j+1), \quad j=0,1,...,n, \tag{6}$$
$$b_j=2(j(j+n-k+3)+n-k+2), \quad j=0,1,...,n-1,$$
$$c_j=4(n-j), \quad j=0,1,...,n-1.$$

2. The other possible substitution is

$$\psi=e^{-\frac{r^4}{8a^4}}e^{-\epsilon\frac{r^2}{4a^2}}r^l e^{-il\phi}w(r), \tag{7}$$

where $l \geq 0$ as before. This gives us the following regular part of solution

$$w(r)=\mathrm{HeunB}(l,\epsilon,-3l+2k,\epsilon l-\lambda,\frac{r^2}{2a^2}). \tag{8}$$

Therefore, $\alpha=l,\ \beta=\epsilon,\ \gamma=-3l+2k, \delta=-\epsilon l-\lambda$. The condition (23) in this case becomes $-2l+k=n+1$. Thus, $k \geq n+1$, the Hamiltonian parameter k must be a natural number and $k-n-1$ must be a non-negative even integer.

Subsequent consideration of this case is the same as in a previous case except for the choice of the permissible $n: n \geq 0,\ n=k-1,k-3,k-5,...$. We choose only real λ in this case as well as in previous one.

The diagonals of the matrix A_{n+1} (22) in this case are defined by the sequences

$$a_j=\lambda-\epsilon(k-n+2j), \quad j=0,1,...,n, \tag{9}$$

$$b_j = j(2j-n+k+3)-n+k+1, \quad j=0,1,...,n-1,$$
$$c_j = 4(n-j), \quad j=0,1,...,n-1.$$

Example 2. Non-rational case

This is defined by the vector potential and potential

$$A_\phi = -\frac{c\hbar}{e}\, k\frac{a}{r\sqrt{r^2+a^2}}, \quad A_r = 0, \quad u = \frac{\hbar^2}{8ma^2}\left(\frac{3a^4}{(r^2+a^2)^2} - \epsilon\frac{a^2}{r^2+a^2}\right), \tag{10}$$

and the energy of the form

$$E = -\frac{\hbar^2}{2ma^2}\chi^2. \tag{11}$$

The magnetic field is equal to

$$B = k\frac{c\hbar}{e}\frac{a}{(r^2+a^2)^{\frac{3}{2}}}.$$

It is interesting that the full magnetic flux through the whole plane is $\Phi = k\frac{2\pi\hbar c}{e}$, that is, it is an even multiple of the quantum flux.

Let us use the change of variables

$$t = \frac{1}{2a}(a+\sqrt{r^2+a^2}). \tag{12}$$

We solve the equation (1) with potentials (10) in order to obtain two linearly independent solutions which can be regular, but not simultaneously, because the condition (30) differs in these cases. We will consider both possibilities separately.

In the first case, we obtain the wave function

$$\psi = e^{il\phi}\sqrt{2t-1}\, t^{\frac{k-l}{2}}(t-1)^{\frac{k+l}{2}} e^{2\chi t}\ \mathrm{HeunC}(4\chi, -l+k, k+l, 0, \frac{1}{2}(k^2-l^2)+\frac{1}{4}(\epsilon+1)-\chi^2, t). \tag{13}$$

Notice, that $k+l$ is a non-negative integer, because the solution must be regular at the point $t=1$ ($r=0$). The condition (30) in this case becomes $n+k+1=0$. It means that k is a negative integer. We substitute the parameters of HeunC function as well as the necessary condition $n=-k-1$ into (29) in order to obtain the diagonals of the matrix A_{n+1} :

$$a_j = \chi(\chi+2(2j-n-l))+l^2-n^2-n-j(j-2n-1)-\frac{1}{4}(1+\epsilon), \quad j=0,1,...,n, \tag{14}$$

$$b_j = (j+1)(j-n-l), \quad j=0,1,...,n-1,$$
$$c_j = 4(n-j)\chi, \quad j=0,1,...,n-1.$$

The function $\det(A_{n+1})$ is a polynomial in χ of degree $2(n+1)$ and the condition $\det(A_{n+1})=0$ defines $2(n+1)$ values of χ. We must choose only negative ones in order to obtain the regular (exponentially decreasing with a distance) wave functions. Only quantum states with the negative χ belong to the discrete spectrum.

In this case we obtain the infinite set of matrices A_{n+1} of the same size $n+1=-k$, where the permissible l are:

$$l = -k, 1-k, 2-k, ...$$

The procedure to obtain the energy levels and wave function is the same as in the previous example except for the definition of the energy levels which are given by (11).

The scale of change of the wave function is $\frac{a}{|\chi|}$ and can be arbitrary a priori.

In the second case, we obtain the wave function in the form

$$\psi = e^{il\phi}\sqrt{2t-1}\,t^{\frac{l-k}{2}}(t-1)^{\frac{k+l}{2}}e^{2\chi t}\ \mathrm{HeunC}(4\chi, l-k, k+l, 0, \frac{1}{2}(k^2-l^2)+\frac{1}{4}(\epsilon+1)-\chi^2, t). \quad (15)$$

We note that $k+l$ is a non-negative integer, because the point $t=1$ ($r=0$) must be regular, as before. The condition (30) in this case takes the form $n+l+1=0$. This means that l must be a negative integer and therefore k must be natural number. We substitute the parameters of HeunC function as well as necessary condition $l=-n-1$ into (29) to obtain the diagonals of the matrix A_{n+1} :

$$a_j = \chi(\chi+2(2j-k-n))-j(j-2n-1)+n+\frac{1}{4}(3-\epsilon), \quad b_j=(j+1)(j-n-k), \quad (16)$$

$$c_j = 4(n-j)\chi.$$

In this case, we obtain a finite set of the matrices A_{n+1} with the permissible n :

$$n \geq 0, \quad n = k-1, k-2, ..., 0.$$

The procedure to obtain the energy levels (by formula (11) in this case) and the wave function is same as in the previous example, as well.

We must choose only negative χ to obtain the quantum states of the discrete spectrum as well as in the previous case.

We denote that values of the discrete spectrum and the corresponding wave functions can be obtained only numerically. We postpone the numerical analysis of this problem for the future work.

Conclusion. In this article, we have considered two absolutely different integrable cases of two-dimensional quantum mechanics with the electromagnetic field. Both cases reveal the rich and complicated structure of the quantum states. These examples were integrated by use of the Confluent and Biconfluent Heun functions. In our opinion, the Heun function and its four confluent forms will be the main special functions of the 21st century. We expect that many problems of quantum mechanics, as well as other applications, will be resolved in terms of these transcendents. However, many results can be obtained only numerically due to the complicated properties of the Heun functions (e.g. calculation of large determinants and solution of the corresponding eigenvalue problems, as in our case).

The approach proposed in this paper is based on the property of quantum Liouville integrability. We hope that it can be applied for a wide range of two-dimensional quantum mechanical problems.

Acknowledgments. The author thanks V.E. Adler, I.V. Kolokolov, and all participants of the mathematical physics seminar at the L.D. landau Institute for Theoretical Physics for the useful discussions. This research was supported in part by the Russian Foundation for Basic Research (Grant No. 16-01-00289).

Appendix

HeunB function

The Biconfluent Heun function satisfies the following equation:

$$\mathrm{BHE} = y''(z) + \left(-2z - \beta + \frac{1+\alpha}{z}\right) y'(z) + \left(\gamma - \alpha - 2 - \frac{1}{2z}((1+\alpha)\beta + \delta)\right) y(z) = 0. \quad (17)$$

Its general solution is

$$y(z) = C_1 \,\mathrm{HeunB}(\alpha, \beta, \gamma, \delta, z) + C_2 \,\mathrm{HeunB}(-\alpha, \beta, \gamma, \delta, z) z^{-\alpha}. \quad (18)$$

Our aim here is to derive the necessary and sufficient conditions for the function $\mathrm{HeunB}(\alpha, \beta, \gamma, \delta, z)$ to be polynomial. To do this in a most convenient way, let us denote

$$\mathrm{HeunB}(\alpha, \beta, \gamma, \delta, z) = P_n(z) = \sum_{j=0}^{n} p_j z^j, \quad (19)$$

then we obtain the following recurrent relation for the coefficients p_j :

$$RB(j) = c_{j-1} p_{j-1} + a_j p_j + b_j p_{j+1} = 0, \quad (20)$$

where

$$a_j = -(\delta + \beta(2j + \alpha + 1)),\ b_j = 2(j(j + \alpha + 2) + \alpha + 1),\ c_j = 2(\gamma - \alpha - 2j - 2). \quad (21)$$

We compute all p_j starting from the initial values $p_{-1} = 0,\ p_0 = 1$. Then, the condition $b_{n+1} = 0$ is equivalent to vanishing of the determinant of the following three-ridiagonal matrix:

$$A_{n+1} = \begin{pmatrix} a_0 & b_0 & 0 & \cdots & 0 \\ c_0 & a_1 & b_1 & \cdots & 0 \\ 0 & c_1 & a_2 & \cdots & \vdots \\ \vdots & \vdots & \vdots & \ddots & b_{n-1} \\ 0 & 0 & \cdots & c_{n-1} & a_n \end{pmatrix}. \quad (22)$$

The second necessary condition can be derived by the following trick. We substitute $y(z) = z^n$ to (17) and calculate the numerator (polynomial) of the obtained expression. The vanishing of the leading term of this polynomial implies the condition $\gamma - \alpha = 2(n+1)$. In this case, if the matrix A_{n+1} is degenerate then $b_{n+1} = 0,\ b_{n+2} = 0$.

Now, we can formulate the following statement.

Statement 1 *The function* $\mathrm{HeunB}(\alpha, \beta, \gamma, \delta, z)$ *equals to a polynomial* $P_n(z)$ *of degree* n *if and only if the following two conditions are satisfied:*

$$\gamma - \alpha = 2(n+1), \quad (23)$$

$$\det(A_{n+1}) = 0. \quad (24)$$

HeunC **function**

The Confluent Heun function satisfies the equation

$$\mathrm{CHE} = y''(z) + \left(\alpha + \frac{\beta+1}{z} + \frac{\gamma+1}{z-1}\right) y'(z) + \left(\frac{\mu}{z} + \frac{\nu}{z-1}\right) y(z) = 0, \tag{25}$$

where $\mu = \frac{1}{2}(\alpha - \beta - \gamma + \alpha\beta - \gamma\beta) - \eta, \quad \nu = \frac{1}{2}(\alpha + \beta + \gamma + \gamma\alpha + \gamma\beta) + \delta + \eta$. Then general solution of this equation is

$$y(z) = C_1\,\mathrm{HeunC}(\alpha, \beta, \gamma, \delta, \eta, z) + C_2\,\mathrm{HeunC}(\alpha, -\beta, \gamma, \delta, \eta, z) z^{-\beta}. \tag{26}$$

Let $\mathrm{HeunC}(\alpha, \beta, \gamma, \delta, z) = P_n(z)$,

$$P_n(z) = \sum_{j=0}^{n} p_j z^j, \tag{27}$$

then we obtain the following recurrent relation for the coefficients p_j :

$$RC(j) = c_{j-1}p_{j-1} + a_j p_j + b_j p_{j+1} = 0, \tag{28}$$

where

$$a_j = \mu - j(j - \alpha + \beta + \gamma + 1), \quad b_j = (j+1)(j+\beta+1), \tag{29}$$
$$c_j = (n-j)\alpha.$$

The same scheme as in the previous section brings us to the following statement.

Statement 2 *The function* $\mathrm{HeunC}(\alpha, \beta, \gamma, \delta, z)$ *equals to a polynomial* $P_n(z)$ *of degree* n *if and only if the following two conditions are satisfied:*

$$\delta = -(n + 1 + \frac{1}{2}(\beta + \gamma))\alpha, \tag{30}$$

$$\det(A_{n+1}) = 0. \tag{31}$$

References

[1] L.D. Landau, E.M. Lifshitz. Vol. 3. Quantum mechanics (non-relativistic theory), third edition, Pergamon, 1991.

[2] L. Infeld, T.E. Hull, The Factorization Method, Reviews of modern physics, **23** (1), 21 (1951).

[3] E. Schrödinger, A method of determining quantum-mechanical eigenvalues and eigenfunctions, Proc. Roy. Irish Acad. **A 46** (1940) 9–16.

[4] E. Schrödinger, Further studies on solving eigenvalue problems by factorization, Proc. Roy. Irish Acad. **A 46** (1941) 183–206.

[5] E. Schrödinger, The factorization of hypergeometric equation, Proc. Roy. Irish Acad. **A 47** (1941) 53–54.

[6] A.M. Ishkhanyan, Exact solution of the Schrödinger equation for the inverse square root potential, Eur. Phys. Lett. **112**, 10006 (2015).

[7] A. Turbiner, Comm. Math. Phys. **118**, 467 (1988).

[8] A. Ushveridze, Sov. Journal Part. Nucl. **20**, 185 (1989).

[9] M.A. Shifman, Int. Journal of Mod. Phys. **A4**, 2897 (1989).

[10] E.V. Ferapontov and A.P. Veselov, J. Math. Phys., 42, 590–607 (2001).

[11] B.A. Dubrovin, I.M. Krichever, and S.P. Novikov, Sov. Math. Dokl., 17, 947–951 (1976).

[12] S.P. Novikov and A.P. Veselov, Amer. Math. Soc. Transl. Ser. 2, 179, 109–132 (1997).

[13] J. Berube and P. Winternitz, J. Math. Phys., 45, 1959–1973 (2004).

[14] K. Heun, Math. Ann., 33, 161 (1889).

[15] H. Bateman and A. Erdélyi, Higher Transcendental Functions, vol 3 (New York, Toronto, London: Mc Grow-Hill Comp. INC, 1955)

[16] A. Decarreau, M.Cl. Dumont-Lepage, P. Maroni, A. Robert and A. Roneaux, Ann. Soc. Bruxelles 92, 53 (1978).

[17] A. Decarreau, P. Maroni and A. Robert, Ann. Soc. Bruxelles 92, 151 (1978).

[18] A. Ronveaux (ed.), Heun's Differential Equations, Oxford Univ. Press, New York (1995).

[19] Plamen P. Fiziev, Novel relations and new properties of confluent Heun's functions and their derivatives of arbitrary order, J Phys. A: Math. Theor. **43** 035203 (2010).

[20] V.G. Marikhin, Quasi-Stäckel systems and two-dimensional Schrödinger equations in an electromagnetic field, Theor. Math. Phys., 177(1), 1352–1360 (2013).